

Package ‘cols’

October 16, 2024

Type Package

Title Constrained Ordinary Least Squares

Version 1.2

Date 2024-10-16

Author Michail Tsagris [aut, cre]

Maintainer Michail Tsagris <mtsagris@uoc.gr>

Depends R (>= 4.0)

Imports quadprog, Rfast, Rfast2

Description Constrained ordinary least squares is performed. One constraint is that all beta coefficients (including the constant) cannot be negative. They can be either 0 or strictly positive. Another constraint is that the sum of the beta coefficients equals a constant. References: Hansen, B. E. (2022). Econometrics, Princeton University Press. <ISBN:9780691235899>.

License GPL (>= 2)

NeedsCompilation no

Repository CRAN

Date/Publication 2024-10-16 11:40:02 UTC

Contents

cols-package	2
Constrained least squares	2
Positive and unit sum constrained least squares	4
Positively constrained least squares	5
Positively constrained least squares with a multivariate response	6

Index	7
--------------	----------

cols-package

Constrained Ordinary Least Squares

Description

Constrained ordinary least squares is performed. One constraint is that all beta coefficients (including the constant) cannot be negative. They can be either 0 or strictly positive. Another constraint is that the sum of the beta coefficients equals a constant. References: Hansen, B.E. (2022). Econometrics, Princeton University Press.

Details

Package: cols
Type: Package
Version: 1.2
Date: 2024-10-16

Maintainers

Michail Tsagris <mtsagris@uoc.gr>.

Author(s)

Michail Tsagris <mtsagris@uoc.gr>

References

Hansen, B. E. (2022). Econometrics, Princeton University Press.

Constrained least squares

Constrained least squares

Description

Constrained least squares.

Usage

```
cls(y, x, R, ca)
mcls(y, x, R, ca)
```

Arguments

y	The response variable. For the <code>cls()</code> a numerical vector with observations, but for the <code>mcls()</code> a numerical matrix .
x	A matrix with independent variables, the design matrix.
R	The R vector that contains the values that will multiply the beta coefficients. See details and examples.
ca	The value of the constraint, $R^T\beta = c$. See details and examples.

Details

This is described in Chapter 8.2 of Hansen (2019). The idea is to minimise the sum of squares of the residuals under the constraint $R^T\beta = c$. As mentioned above, be careful with the input you give in the x matrix and the R vector. The `cls()` function performs a single regression model, whereas the `mcls()` function performs a regression for each column of y. Each regression is independent of the others.

Value

A list including:

be	A numerical matrix with the constrained beta coefficients.
mse	A numerical vector with the mean squared error.

Author(s)

Michail Tsagris.

R implementation and documentation: Michail Tsagris <mtsagris@uoc.gr>.

References

Hansen, B. E. (2022). *Econometrics*, Princeton University Press.

See Also

[pls](#)

Examples

```
x <- as.matrix( iris[1:50, 1:4] )
y <- rnorm(50)
R <- c(1, 1, 1, 1)
cls(y, x, R, 1)
```

Positive and unit sum constrained least squares

Positive and unit sum constrained least squares

Description

Positive and unit sum constrained least squares.

Usage

```
pcls(y, x)
mpcls(y, x)
```

Arguments

y	The response variable. For the pcls() a numerical vector with observations, but for the mpcls() a numerical matrix.
x	A matrix with independent variables, the design matrix.

Details

The constraint is that all beta coefficients are positive and sum to 1. The pcls() function performs a single regression model, whereas the mpcls() function performs a regression for each column of y. Each regression is independent of the others.

Value

A list including:

be	A numerical matrix with the positively constrained beta coefficients.
mse	A numerical vector with the mean squared error.

Author(s)

Michail Tsagris.

R implementation and documentation: Michail Tsagris <mtsagris@uoc.gr>.

See Also

[pls](#), [cls](#), [mvpls](#)

Examples

```
x <- as.matrix( iris[1:50, 1:4] )
y <- rnorm(50)
pcls(y, x)
```

Positively constrained least squares

Positively constrained least squares

Description

Positively constrained least squares.

Usage

```
pls(y, x)
mpls(y, x)
```

Arguments

y	The response variable. For the pls() a numerical vector with observations, but for the mpls() a numerical matrix .
x	A matrix with independent variables, the design matrix.

Details

The constraint is that all beta coefficients (including the constant) are non negative. The pls() function performs a single regression model, whereas the mpls() function performs a regression for each column of y. Each regression is independent of the others.

Value

A list including:

be	A numerical matrix with the positively constrained beta coefficients.
mse	A numerical vector with the mean squared error(s).

Author(s)

Michail Tsagris.

R implementation and documentation: Michail Tsagris <mtsagris@uoc.gr>.

See Also

[cls](#), [pcls](#), [mvpls](#)

Examples

```
x <- as.matrix( iris[1:50, 1:4] )
y <- rnorm(50)
pls(y, x)
```

Positively constrained least squares with a multivariate response

Positively constrained least squares with a multivariate response

Description

Positively constrained least squares with a multivariate response.

Usage

```
mvpls(y, x)
```

Arguments

y	The response variables, a numerical matrix with observations.
x	A matrix with independent variables, the design matrix.

Details

The constraint is that all beta coefficients (including the constant) are positive.

Value

A list including:

be	The positively constrained beta coefficients.
mse	The mean squared error.

Author(s)

Michail Tsagris.

R implementation and documentation: Michail Tsagris <mtsagris@uoc.gr>.

See Also

[cls](#)

Examples

```
y <- as.matrix( iris[, 1:2] )
x <- as.matrix( iris[, 3:4] )
mvpls(y, x)
```

Index

cls, [4-6](#)
cls (Constrained least squares), [2](#)
cols-package, [2](#)
Constrained least squares, [2](#)

mcls (Constrained least squares), [2](#)
mpcls (Positive and unit sum
constrained least squares), [4](#)
mpls (Positively constrained least
squares), [5](#)
mvpls, [4, 5](#)
mvpls (Positively constrained least
squares with a multivariate
response), [6](#)

pcls, [5](#)
pcls (Positive and unit sum
constrained least squares), [4](#)
pls, [3, 4](#)
pls (Positively constrained least
squares), [5](#)
Positive and unit sum constrained
least squares, [4](#)
Positively constrained least squares, [5](#)
Positively constrained least squares
with a multivariate response, [6](#)