

EUROPEAN ORGANIZATION FOR NUCLEAR RESEARCH

CERN-PH/TN/2005-49  
30 February 2005

**TEST-BEAM PERFORMANCE OF A TRACKING TRD PROTOTYPE**

**Abstract**

Each paper should be preceded by a short Abstract of not more than 150 words, which should be written as a single paragraph and should not contain references.

J.T. Shank, J.S. Whitaker  
Boston University, Boston, Mass., USA.

V.A. Polychronakos, V. Radeka, D. Stephani  
Brookhaven National Laboratory, Upton, NY, USA.

H. Beker, R.K. Bock, M. Botlo, L. DiCiaccio<sup>c)</sup>, M. Koratzinos, A. Lopez<sup>\*</sup>)  
CERN, Geneva, Switzerland.

W.J. Willis  
Columbia University, New York, NY, USA.

T. Akesson  
Lund University, Lund, Sweden.

V. Chernyatin, B. Dolgoshein, P. Nevski  
Moscow Physical Engineering Institute, Moscow, USSR.

*Our collaboration*

*(Submitted to some important Journal)*

*To be presented by the author  
on the conference in somewhere at sometimes.*

*NOTE: not yet final*

*Dedicated to AAA*

---

<sup>c)</sup>Univ. di Roma II, ‘Tor Vergata’, Rome, Italy.

<sup>\*</sup>)On leave from Fac. de Ciencias Físicas, Univ. Complutense, Madrid, Spain.

## 1 INTRODUCTION

The biological effect of radiation depends on the quality of the radiation as well as on the amount of energy absorbed. Evidence suggests that this quality dependence is primarily caused by the differences in rates of energy loss [1, 2]. The general criteria that have been used are:

- i) to terminate the iteration when the residue between iterated and experiment values is of the order of experimental errors [1, 3–6];
- ii) to terminate when the smoothest solution has been obtained.

For an overall appreciation of the work carried out in this field, see Refs. [7] and [8].

## 2 RADIATION PARAMETERS EFFECTIVE IN CAUSING BIOLOGICAL DAMAGE

### 2.1 Linear energy transfer

Smith tabulated the average values of mean linear energy transfer (LET) obtained by the different methods used in Ref. [3], and these are reproduced in Table 1.

Table 1: Calculated mean LET values in water (keV/mm)

Radiation	Smith <sup>a)</sup>	Jones
1 MeV $\alpha$		
200 kVp X-rays total	3.25	1.79
200 kVp X-rays (primary)	2.60	1.48

<sup>a)</sup> J. Smith, Rad. Res. 1 (1956) 234

Figure 1 is reproduced from the publication mentioned earlier and shows the good agreement between predictions and calculations. Comparison should be made with the decay curves shown in Figs. 6 and 7 of Ref. [4], and further information is given in Section 2.1.2 and Appendix APPENDIX A::.

In order to understand the process more precisely, we must first explain the angle and curvature generation systems.




Fig. 1: Diagram of a straight line.

### 2.1.1 Angle generation

Equation 1 representing a straight line at an angle  $\theta$ , is

$$n^k(h)u = \lambda h t g q_k \quad (1)$$

and

$$n^k(h) = kh \frac{k}{32} \quad (2)$$

where:

$\lambda$  is the distance between two consecutive sweep lines,

$u$  is the least count.

### 2.1.2 Curvature generation

We consider a parabola, the tangent of which is parallel to the vertical axis of the main matrix. A sufficiently good approximation to a parabola [see Eq. 3] drawn inside the matrix can be given by the following formula:

$$n_q = \alpha q^2 \quad \alpha = \frac{\lambda^2}{3Ru} \quad (3)$$

where  $R$  is the radius of curvature. We have the following relationship:

$$n_q + \mu_q \text{ with } \mu_q = \alpha(2q + 1) \quad (4)$$

## 3 CONCLUSION

The theoretical considerations presented have been confirmed by their close agreement with the results of practical experiments. An account of the earlier work carried out in this field can be found in the bibliography.

It is expected that in the next few years many new results will be published, since a significant number of new experiments have recently been launched.

## **ACKNOWLEDGEMENTS**

We wish to thank C. Brown for his most enlightening comments on this topic.

## **REFERENCES**

- [1] J.M. Raby, Biophysical aspects of radiation quality, International Atomic Energy Agency, Technical Reports Series No. 58 (1966).
- [2] J.-P. Dupont, Proc. Int. Conf. on Radiation Hazards, Columbia, 1960 (Academic Press Inc., New York, 1961), Vol. II, p. 396.
- [3] H. Appleman et al., *J. Med. Biol.* **8** (1959) 911.
- [4] E. van Berg, D. Johnson and J. Smith, *Rad. Res.* **5** (1965) 215.
- [5] P. Bryant and S. Newman (eds.), The generation of high fields, CAS-ECFA-INFN Workshop, Frascati, 1984. (ECFA 85/91, CERN 85/07, CERN, Geneva, 1985).
- [6] M.A. Allen et al., *IEEE Trans. Nucl. Sci.* **NS-24** (1977) 1780.
- [7] E. Keil, *Nucl. Instrum. Methods* **100** (1969) 419.
- [8] G. Guignard, CERN LEP-TH/83-38 (1983).

## **BIBLIOGRAPHY**

I.C. Percival and D. Richards, Introduction to dynamics (Cambridge University Press, 1982).  
L. Garrido (ed.), Dynamical systems and chaos: Proc. Sitges 1982, Lecture Notes in Physics No. 179 (Springer, Berlin, 1983).

## **APPENDIX A: CONSTRUCTION ON A FLAT SITE**

### **A.1 General considerations**

Following on ECFA recommendation, the project described in this report is based on the assumption that the machine is built close to the present CERN site, and has been taken into account in the cost estimate.

### **A.2 Effects on the construction**

The general layout of the machine would be very similar to that shown in the main body of the report.